

# QUESTIONS TO ASK YOUR HEALTHCARE PROVIDER

When you meet with your healthcare provider, it's important that you discuss any concerns or questions you may have about your disease and your treatment for it. To help you start the conversation, here are some questions for you to consider asking. If you have additional questions or perhaps topics you would like to know more about, it's a good idea to write them down and add them to this list whenever you think of any. Bring these pages with you to your next appointment. Remember to jot down the answers you receive, so you can refer to them later.

WHAT STAGE OF MELANOMA DO I HAVE?

---

---

---

HAS MY MELANOMA SPREAD TO OTHER PARTS OF MY BODY?

---

---

---

HOW LONG DO I NEED TO TAKE BRAFTOVI + MEKTOVI?

---

---

---

ARE THERE ANY SYMPTOMS OR SIDE EFFECTS I SHOULD TELL YOU ABOUT RIGHT AWAY?

---

---

---

**Please see following pages for Important Safety Information, full BRAFTOVI Prescribing Information, including Medication Guide, and full MEKTOVI Prescribing Information, including Patient Information, for additional information.**

WHAT CAN I DO TO HELP MANAGE SIDE EFFECTS?

---

---

---

AM I GOING TO LOSE MY HAIR?

---

---

---

WHAT SHOULD MY FAMILY KNOW ABOUT MY TAKING BRAFTOVI + MEKTOVI?

---

---

---

SHOULD I STILL GO FOR SKIN EXAMS? HOW OFTEN?

---

---

---

WILL I NEED TO CHANGE MY DOSE OF BRAFTOVI OR MEKTOVI DURING TREATMENT?

---

---

---

**Please see following pages for Important Safety Information, full BRAFTOVI Prescribing Information, including Medication Guide, and full MEKTOVI Prescribing Information, including Patient Information, for additional information.**

## INDICATION AND IMPORTANT SAFETY INFORMATION

### What is BRAFTOVI + MEKTOVI?

BRAFTOVI and MEKTOVI are prescription medicines used together to treat people with a type of skin cancer called melanoma:

- that has spread to other parts of the body or cannot be removed by surgery, **and**
- that has a certain type of abnormal “BRAF” gene

BRAFTOVI should not be used to treat people with wild-type BRAF melanoma. Your healthcare provider will perform a test to make sure that BRAFTOVI + MEKTOVI is right for you. It is not known if BRAFTOVI or MEKTOVI is safe and effective in children.

### IMPORTANT SAFETY INFORMATION

#### What is the most important information I should know about BRAFTOVI?

##### **BRAFTOVI may cause serious side effects, including:**

- **Risk of new skin cancers.** BRAFTOVI, when used alone or with MEKTOVI, may cause skin cancers called cutaneous squamous cell carcinoma or basal cell carcinoma

Talk to your healthcare provider about your risk for these cancers

Check your skin and tell your healthcare provider right away about any skin changes, including a new wart, skin sore or reddish bump that bleeds or does not heal, or change in size or color of a mole

Your healthcare provider should check your skin before and during treatment, and for up to 6 months after you stop treatment to look for any new skin cancers

Your healthcare provider should also check for cancers that may not occur on the skin. Tell your healthcare provider about any new symptoms that develop during treatment

- **Bleeding problems.** BRAFTOVI, when taken with MEKTOVI, can cause serious bleeding problems, including in your stomach or brain, that can lead to death. Call your healthcare provider and get medical help right away if you have any signs of bleeding, including: headaches, dizziness, or feeling weak; cough up blood or blood clots; vomit blood or your vomit looks like “coffee grounds”; red or black stools that look like tar

- **Eye problems.** Tell your healthcare provider right away if you develop any of these symptoms of eye problems: blurred vision, loss of vision, or other vision changes; see colored dots; see halos (blurred outline around objects); eye pain, swelling, or redness
- **Changes in the electrical activity of your heart called QT prolongation.** QT prolongation can cause irregular heartbeats that can be life-threatening. Your healthcare provider should do tests before you start and during your treatment to check your body salts (electrolytes). Tell your healthcare provider right away if you feel faint, light-headed, dizzy, or if you feel your heart beating irregularly or fast

##### **MEKTOVI may cause serious side effects, including:**

- **Heart problems, including heart failure.** Your healthcare provider should check your heart function before and during treatment. Call your healthcare provider right away if you have any of the following signs and symptoms of a heart problem: feeling like your heart is pounding or racing; shortness of breath; swelling of your ankles and feet; feeling light-headed
- **Blood clots.** MEKTOVI can cause blood clots in your arms or legs, which can travel to your lungs and can lead to death. Get medical help right away if you have the following symptoms: chest pain; sudden shortness of breath or trouble breathing; pain in your legs with or without swelling; swelling in your arms and legs; a cool, pale arm or leg
- **Eye problems.** MEKTOVI can cause serious eye problems that might lead to blindness. Call your healthcare provider right away if you develop any of these symptoms of eye problems: blurred vision, loss of vision, or other vision changes; see colored dots; see halos (blurred outline around objects); eye pain, swelling, or redness
- **Lung or breathing problems.** MEKTOVI can cause lung or breathing problems. Tell your healthcare provider if you have any new or worsening symptoms of lung or breathing problems, including: shortness of breath; cough

**Please see following pages for full BRAFTOVI Prescribing Information, including Medication Guide, and full MEKTOVI Prescribing Information, including Patient Information, for additional information.**

  
**BRAFTOVI™ + MEKTOVI®**  
(encorafenib) capsules      (binimetinib) tablets

- **Liver problems.** Your healthcare provider should perform blood tests to check your liver function before and during treatment. Tell your healthcare provider if you have any of the following signs and symptoms of a liver problem: yellowing of your skin or the white part of your eyes (jaundice); dark or brown (tea-colored) urine; nausea or vomiting; loss of appetite
  - **Muscle problems (rhabdomyolysis).** MEKTOVI can cause muscle problems that can be severe. Your healthcare provider should perform a blood test to check your levels of an enzyme in your blood called creatine phosphokinase (CPK) before and during treatment. Tell your healthcare provider right away if you develop any of these symptoms: weakness; muscle aches or pain; dark, reddish urine
  - **Bleeding problems.** MEKTOVI, when taken with BRAFTOVI, can cause serious bleeding problems, including in your brain or stomach, that can lead to death. Call your healthcare provider and get medical help right away if you have any signs of bleeding, including: headaches, dizziness, or feeling weak; cough up blood or blood clots; vomit blood or your vomit looks like “coffee grounds”; red or black stools that look like tar
- have heart problems, including a condition called long QT syndrome
  - have been told that you have low blood levels of potassium, calcium, or magnesium
  - have liver or kidney problems
  - have had blood clots
  - have lung or breathing problems
  - have any muscle problems
  - have high blood pressure (hypertension)
  - are pregnant or plan to become pregnant. BRAFTOVI + MEKTOVI can harm your unborn baby.
    - Females who are able to become pregnant should use effective nonhormonal birth control (contraception) during treatment and for at least 30 days after the final dose. Birth control methods that contain hormones (such as birth control pills, injections, or transdermal systems) may not work as well during treatment with BRAFTOVI
    - Tell your healthcare provider right away if you become pregnant or think you might be pregnant during treatment with BRAFTOVI + MEKTOVI
  - are breastfeeding or plan to breastfeed. Do not breastfeed during treatment and for 2 weeks after the final dose. Talk to your healthcare provider about the best way to feed your baby during this time

**The most common side effects of BRAFTOVI and MEKTOVI when taken in combination:** fatigue, nausea, diarrhea, vomiting, pain or swelling of your joints, abdominal pain

BRAFTOVI may cause fertility problems in males. Talk to your healthcare provider if this is a concern for you.

**These are not all the possible side effects of BRAFTOVI and MEKTOVI.**

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Array BioPharma Inc. at 1-844-792-7729.**

**Before taking BRAFTOVI + MEKTOVI, tell your healthcare provider about all of your medical conditions, including if you:**

- have had bleeding problems
- have eye problems

**Please see following pages for full BRAFTOVI Prescribing Information, including Medication Guide, and full MEKTOVI Prescribing Information, including Patient Information, for additional information.**



MEKTOVI® is a registered trademark of Array BioPharma Inc. in the United States and various other countries.

BRAFTOVI™ is a trademark of Array BioPharma Inc.

© 2018 Array BioPharma Inc. All rights reserved. PM-US-BIN+ENC-0141 07/18 vi